Subpart G—Source Plasma

- 640.60 Source Plasma.
- 640.61 Informed consent.
- 640.62 Medical supervision.
- 640.63 Suitability of donor.
- 640.64 Collection of blood for Source Plasma.
- 640.65 Plasmapheresis.
- 640.66 Immunization of donors.
- 640.67 Laboratory tests.
- 640.68 Processing.
- 640.69 General requirements.
- 640.70 Labeling.
- 640.71 Manufacturing responsibility.
- 640.72 Records.
- 640.72 Reporting of fatal donor reactions.
- 640.74 Modification of Source Plasma.
- 640.76 Products stored or shipped at unacceptable temperatures.

Subpart H—Albumin (Human)

- 640.80 Albumin (Human).
- 640.81 Processing.
- 640.82 Tests on final product.
- 640.83 General requirements.
- 640.84 Labeling.

Subpart I—Plasma Protein Fraction (Human)

- 640.90 Plasma Protein Fraction (Human).
- 640.91 Processing.
- 640.92 Tests on final product.
- 640.93 General requirements.
- 640.94 Labeling.

Subpart J—Immune Globulin (Human)

- 640.100 Immune Globulin (Human).
- 640.101 General requirements.
- 640.102 Manufacture of Immune Globulin (Human).
- 640.103 The final product.
- 640.104 Potency.

Subpart K [Reserved]

Subpart L—Alternative Procedures

640.120 Alternative procedures.

AUTHORITY: 21 U.S.C. 321, 351, 352, 353, 355, 360, 371; 42 U.S.C. 216, 262, 263, 263a, 264.

Source: $38\ FR\ 32089,\ Nov.\ 20,\ 1973,\ unless$ otherwise noted.

CROSS REFERENCES: For U.S. Customs Service regulations relating to viruses, serums, and toxins, see 19 CFR 12.21–12.23. For U.S. Postal Service regulations relating to the admissibility to the United States mails see parts 124 and 125 of the Domestic Mail Manual, that is incorporated by reference in 39 CFR part 111.

Subpart A—Whole Blood

§ 640.1 Whole Blood.

The proper name of this product shall be Whole Blood. Whole Blood is defined as blood collected from human donors for transfusion to human recipients.

[38 FR 32089, Nov. 20, 1973, as amended at 50 FR 4138, Jan. 29, 1985]

§ 640.2 General requirements.

- (a) Manufacturing responsibility. All manufacturing of Whole Blood, including donor examination, blood collection, laboratory tests, labeling, storage and issue, shall be done under the supervision and control of the same licensed establishment except that the Director, Center for Biologics Evaluation and Research, may approve arrangements, upon joint request of two or more licensed establishments, which he finds are of such a nature as to assure compliance otherwise with the provisions of this subchapter.
- (b) Blood container. The blood container shall not be entered prior to issue for any purpose except for blood collection or when the method of processing requires use of a different concontainer shall The tainer. uncolored and transparent to permit visual inspection of the contents and any closure shall be such as will maintain an hermetic seal and prevent contamination of the contents. The container material shall not interact with the contents under the customary conditions of storage and use, in such a manner as to have an adverse effect upon the safety, purity, or potency of the blood.
- (c) *Reissue of blood*. Blood that has been removed from storage controlled by a licensed establishment shall not be reissued by a licensed establishment unless the following conditions are observed:
- (1) The container has a tamper-proof seal when originally issued and this seal remains unbroken;
- (2) A segment is properly attached and has not been removed, except that blood lacking a properly attached segment may be reissued in an emergency provided it is accompanied by instructions for sampling and for use within 6

§ 640.3

hours after entering the container for sampling;

- (3) The blood has been stored continuously at 1 to 6 °C and shipped between 1 and 10 °C;
- (4) The blood is held for observation until a significant inspection consistent with the requirements of §640.5(e) can be made.

[38 FR 32089, Nov. 20, 1973, as amended at 41 FR 4015, Jan. 28, 1976; 42 FR 59878, Nov. 22, 1977; 43 FR 34460, Aug. 4, 1978; 49 FR 15187, Apr. 18, 1984; 49 FR 23834, June 8, 1984; 50 FR 4138, Jan. 29, 1985; 53 FR 116, Jan. 5, 1988; 55 FR 11013, Mar. 26, 1990; 63 FR 16685, Apr. 6, 1998; 64 FR 45371, Aug. 19, 1999; 66 FR 1836, Jan. 10, 2001; 66 FR 31165, June 11, 2001; 66 FR 40889, Aug. 6, 2001; 67 FR 9587, Mar. 4, 2002]

§ 640.3 Suitability of donor.

- (a) Method of determining. The suitability of a donor as a source of Whole Blood shall be determined by a qualified physician or by persons under his supervision and trained in determining suitability. Such determination shall be made on the day of collection from the donor by means of medical history, a test for hemoglobin level, and such physical examination as appears necessary to a physician who shall be present on the premises when examinations are made, except that the suitability of donors may be determined when a physician is not present on the premises, provided the establishment (1) maintains on the premises, and files with the Center for Biologics Evaluation and Research, a manual of standard procedures and methods, approved by the Director of the Center for Biologics Evaluation and Research, that shall be followed by employees who determine suitability of donors, and (2) maintains records indicating the name and qualifications of the person immediately in charge of the employees who determine the suitability of donors when a physician is not present on the premises.
- (b) Qualifications of donor; general. Except as provided in paragraph (f) of this section and for autologous donations, a person may not serve as a source of Whole Blood more than once in 8 weeks. In addition, donors shall be in good health, as indicated in part by:
 - (1) Normal temperature;
- (2) Demonstration that systolic and diastolic blood pressures are within

- normal limits, unless the examining physician is satisfied that an individual with blood pressures outside these limits is an otherwise qualified donor under the provisions of this section:
- (3) For allogeneic donors, a blood hemoglobin level which shall be demonstrated to be no less than 12.5 grams (g) of hemoglobin per 100 milliliters (mL) of blood; or a hematocrit value of 38 percent, and for autologous donors, a blood hemoglobin level which shall be demonstrated to be no less than 11.0 g of hemoglobin per 100 mL of blood or a hematocrit value of 33 percent.
- (4) Freedom from acute respiratory diseases:
- (5) Freedom from any infectious skin disease at the site of phlebotomy and from any such disease generalized to such an extent as to create a risk of contamination of the blood;
- (6) Freedom from any disease transmissible by blood transfusion, insofar as can be determined by history and examinations indicated above; and
- (7) Freedom of the arms and forearms from skin punctures or scars indicative of addiction to self-injected narcotics.
- (c) Additional qualifications of donor; viral hepatitis. No individual shall be used as a source of Whole Blood if he
- (1) A history of viral hepatitis after the 11th birthday;
- (2) A history of close contact within 12 months of donation with an individual having viral hepatitis;
- (3) A history of having received within 12 months of donation, human blood or any derivative of human blood which the Food and Drug Administration has advised the blood establishment is a possible source of viral hepatitis.
- (d) Therapeutic bleedings. Blood withdrawn in order to promote the health of a donor otherwise qualified under the provisions of this section, shall not be used as a source of Whole Blood unless the container label conspicuously indicates the donor's disease that necessitated withdrawal of blood.
 - (e) [Reserved]
- (f) Qualifications; donations within less than 8 weeks. A person may serve as a source of Whole Blood more than once